

Q	Answer	Mark	Comments
1(a)	(0).35	B1	oe
	Additional Guidance		
	Mark the answer line. If this is blank, mark the working		
	If values are given in one or more forms, either on the answer line or in working with nothing on the answer line, all values must be correct		
	eg1 $0.35 = \frac{7}{20}$ on answer line	B1	
	eg2 $\frac{35}{100}$ and 3.5 in working with $\frac{35}{100}$ on answer line	B1	
	eg3 $\frac{35}{100}$ and 3.5 in working with 3.5 on answer line	B0	
	eg4 $\frac{35}{100}$ and 3.5 in working with answer line blank	B0	

Q	Answer	Mark	Comments
1(b)	$\frac{5}{18}$	B1	oe eg $\frac{10}{36}$
	Additional Guidance		
	Mark the answer line. If this is blank, mark the working		
	Allow 0.277... (minimum two 7s and two dots) or correct notation for recurring decimals		
	If values are given in one or more forms, either on the answer line or in working with nothing on the answer line, all values must be correct		
	eg1 $\frac{5}{18} = 0.277\dots$ on answer line	B1	
	eg2 $\frac{5}{18}$ and 0.518 in working with answer line blank	B0	
	$\frac{1\frac{2}{3}}{6}$ or $\frac{1.66\dots}{6}$ without answer in correct form	B0	

Q	Answer	Mark	Comments
1(c)	45	B1	
	Additional Guidance		
	Mark the answer line. If this is blank, mark the working		
	If values are given in one or more forms, either on the answer line or in working with nothing on the answer line, all values must be correct		B1
	eg1 $\frac{270}{6} = 45$ on answer line		B0
	eg2 $\frac{270}{6}$ and $44\frac{5}{6}$ in working with answer line blank		B0
	Do not allow unprocessed answers		B0
	eg $\frac{270}{6}$		

Q	Answer	Mark	Comments
2	$x < 13$ or $13 > x$	B1	
	Additional Guidance		
	$x = 13$ in working with $x < 13$ on answer line		B1
	$x < 13$ and $(x =) 13$ on answer line		B0
	$x < 13$ in working with $x = 13$ or 13 on answer line		B0
	Ignore number lines drawn		

Q	Answer	Mark	Comments
3	$2\frac{1}{4}$	B1	oe mixed number
	Additional Guidance		
	$\frac{9}{4} = 2\frac{1}{4}$ or $2.25 = 2\frac{1}{4}$ on answer line	B1	
	$2\frac{1}{4} = \frac{9}{4}$ or $2\frac{1}{4} = 2.25$ on answer line	B0	
	Otherwise, $2\frac{1}{4}$ and $\frac{9}{4}$ or $2\frac{1}{4}$ and 2.25 on answer line in either order (or in working with answer line blank and answer unclear)	B0	
	$1\frac{5}{4}$	B0	
	$2\left(\frac{1}{4}\right)$ or $2 + \frac{1}{4}$	B0	

Q	Answer	Mark	Comments
4	Alternative method 1 – numerical		
	1 and 5 and 3 or 9 (parts) or numbers in the ratio 1 : 5 : 3 or (angle sum on a straight line =) 180	M1	oe may be seen in a ratio eg $\frac{1}{5} : 1 : \frac{3}{5}$ or $\frac{1}{3} : \frac{5}{3} : 1$ numbers can be in any order eg 30, 10, 50
	180 \div (1 + 5 + 3) or 20 or $180 \div \frac{9}{5}$	M1dep	oe
	100	A1	
	Alternative method 2 – algebraic		
	x and $5x$ and $3x$ or $9x$ or (angle sum on a straight line =) 180	M1	oe correct terms with any angle as x any letter, any order may be seen on diagram
	Correct equation with correct method to solve for one angle	M1dep	eg $x + 5x + 3x = 180$ and $180 \div (1 + 5 + 3)$
	100	A1	
	Additional Guidance		
	$x + 5x + 3x = 360$ or $360 \div 9$		M1M0A0
	$\frac{1}{5}x + x + \frac{3}{5}x = 180$ and $180 \div \left(\frac{1}{5} + 1 + \frac{3}{5}\right)$		M1M1
	$\frac{1}{3}x + \frac{5}{3}x + x = 180$ and $180 \div \left(\frac{1}{3} + \frac{5}{3} + 1\right)$		M1M1
	Angle EBD marked as 100 on the diagram with answer line blank		M1M1A1
	20 and 100 in working with no or incorrect answer chosen		M1M1A0

Q	Answer	Mark	Comments
5	All conditions met: <ul style="list-style-type: none"> first number is prime second number is prime correctly evaluated even answer answer in range 	B3	if their product is incorrectly evaluated or missing, then 'even answer' and 'answer in range' refer to the correct product for their multiplication B2 4 conditions met B1 3 conditions met
	Additional Guidance $2 \times 29 = 58$ (or $29 \times 2 = 58$) is the only fully correct solution Allow 50 to 60 inclusive for 'answer in range' Award the best mark from boxes or in working for up to B2 The two prime numbers do not have to be different		B3
6			
M1	oe eg $96 \div 6 \times 5$ implied by 176		
M1dep	oe eg $80 \div 4$		
M1	oe eg $96 \div 3 \times 2$ accept 0.66 or better for $\frac{2}{3}$		
A1	SC2 100.8(0) or [77.32, 77.34] condone incorrect money notation eg 84.0 or 84.00p		
6	Additional Guidance SC2 for 100.8(0) is from misreading as Andrew gets £96 SC2 for [77.32, 77.34] is from $\frac{2}{3}$ of 80 plus $\frac{1}{4}$ of 96 Do not accept ' $\frac{5}{6}$ of 96' or ' $\frac{1}{4}$ of 80' or ' $\frac{2}{3}$ of 96' for M marks unless accompanied by a correct method or value		

Q	Answer	Mark	Comments	
7	Alternative method 1 – evaluation and division			
	$(5^2 =) 25$ or $(3 \times 5^2 =) 75$ or $600 \div 3$ or 200 or $600 \div 5^2$ or 24	M1	oe oe eg $3 \times 200 = 600$ oe eg $25 \times 24 = 600$	
	$600 \div 3 \div 5^2$ or 8		M1dep oe eg $8 \times 75 = 600$	
	3 with M1 awarded and not from incorrect working	A1		
	Alternative method 2 – product of prime factors			
	600 written as a product of factors where at least one factor is prime	M1	eg 2 and 300 or 5 and 120 or 2 and 2 and 150 may be seen on a factor tree or in repeated division allow one strand to be incorrect if a previous value completes the product eg 20×30 followed by $2 \times 10 \times 5 \times 8$ implies $2 \times 10 \times 30$ for M1	
	2 and 2 and 2 and 3 and 5 and 5		M1dep may be seen on a factor tree or in repeated division	
	3 with M1 awarded and not from incorrect working	A1		
	Additional Guidance			
	8 \times 3 \times 25 = 600 and answer 3		M1M1A1	
	2 ³ on answer line with M2 awarded		M1M1A0	
	Answer 3 on answer line with no working		M0M0A0	
	Do not allow $600 \div 3 \times 5^2$ for M2 in alt 1 unless recovered, but do allow $\frac{600}{3 \times 5^2}$ or $600 \div (3 \times 5^2)$			

Q	Answer	Mark	Comments
	$13x + 22$	B2	B1 $15x + 20$ or $-2x + 2$ or $13x + a$ or $bx + 22$, where a and b can be any numbers
Additional Guidance			
8	Do not ignore further working for B2 eg $13x + 22 = 35x$ eg $13x + 22, x = \frac{22}{13}$		B1 B1

Q	Answer	Mark	Comments
9	Any two from: Reference to graph passing through point where $x = 0$ Reference to graph being incorrect for negative x values Reference to the graph stopping before the end of the axes/axis	B2	B1 any one correct reference eg the graph touches the y -axis eg the graph to the left of the y -axis should be below the x -axis eg the graph should go to the ends of the axes
Additional Guidance			
	Ignore non-contradictory, irrelevant responses alongside a correct response		
	Draws correct graph		
	Draws graph with one section correct for positive values of x or negative values of x		
	'It isn't the graph of $y = \frac{1}{x}$ ', scores B0, but B1 may still be scored for the other criticism		
	'There are no numbers on the axes' scores B0, but B1 may still be scored for the other criticism		
Mark for graph touching y-axis			
	You cannot have $x = 0$	B1	
	The line in the top right should be moved to the right	B1	
	It says $x \neq 0$ but it (the sketch) does	B1	
	One line is touching the y -axis	B1	
	The lines should be symmetrical	B0	
	You cannot have $y = 0$	B0	
	One line is touching the y -axis but the other isn't	B0	

Question 9 Additional Guidance continues on the next page

9 cont	Mark for negative values being in the wrong quadrant	
	There shouldn't be anything in the top-left section	B1
	There should be something in the bottom-left section	B1
	It is the graph of $y = \frac{1}{x^2}$	B1
	It should have rotational symmetry	B1
	It should be symmetrical about $y = x$	B1
	It should be symmetrical about $y = -x$	B1
	It should be symmetrical	B0
	One should be negative	B0
	The bit on the left is wrong	B0
	The negative values are plotted incorrectly	B0
	Reference to the graph stopping before the end of the axes	
	It stops before the end of the axes	B1
	The lines don't go far enough	B1
	The lines need to be higher up	B0

Q	Answer	Mark	Comments
10	Alternative method 1 – algebra based on Sunita's age		
	5 × 3 or 15	M1	may be implied by their algebraic total of the three ages being divided by 3
	$x - 1$ or $2x$ or $4x - 1$	M1	oe expressions any letter throughout
	$x + \text{their } (x - 1) + \text{their } 2x = \text{their } 15$ or $4x - 1 = \text{their } 15$	M1dep	oe equation eg $\frac{x + x - 1 + 2x}{3} = 5$ dep on M1M1
	$(x =) 4$	M1dep	correct solution to their equation if the solution has a decimal part allow truncation or rounding to the nearest whole number
	8	A1	
	Alternative method 2 – algebra based on Joel's age		
	5 × 3 or 15	M1	may be implied by their algebraic total of the three ages being divided by 3
	$\frac{y}{2}$ or $\frac{y}{2} - 1$ or $2y - 1$	M1	oe expressions any letter throughout $2y - 1$ must not come from $y + y - 1$
	$y + \text{their } \frac{y}{2} + \text{their } (\frac{y}{2} - 1) = \text{their } 15$	M1dep	oe equation eg $\frac{y + \frac{y}{2} + \frac{y}{2} - 1}{3} = 5$ dep on M1M1
	$2y + \text{their } y + \text{their } (y - 2) = 2 \times \text{their } 15$ or $4y - 2 = 30$ or $2y - 1 = 15$	M1dep	their equation with no denominator
	8	A1	

Question 10 continues on the next page

10 cont	Alternative method 3 – trial and improvement		
	5 × 3 or 15	M1	may be implied by their total of the three ages being divided by 3
	Trial of three numbers which fit the criteria, with either their sum correctly evaluated or their sum divided by 3	M1	eg $2 + 1 + 4 = 7$ or $(2 + 1 + 4) \div 3$ condone missing brackets
	Second trial of three numbers which fit the criteria, with either their sum correctly evaluated or their sum divided by 3	M1dep	dep on previous M1 eg $3 + 2 + 6 = 11$ or $(3 + 2 + 6) \div 3$ condone missing brackets
	4, 3 and 8 selected as their final combination	M1dep	any order implies M4
	8	A1	
	Additional Guidance		
	Up to M4 may be awarded for correct work seen in multiple attempts even if not subsequently used		
	Correct expressions, but the sum of the three ages is equated to 5 eg $4x - 1 = 5$		M0M1M0M0A0
	In alt 1, the correct value of x or the correct age for Joel for their two terms for Beth and Joel, with one correct, implies the first 4 marks eg x and $x + 1$ and $2x$, with $x = 3.5$ or answer 7		M1M1M1M1A0
	In alt 2, the correct value of y for their two terms for Sunita and Beth, with one correct, implies the first 4 marks eg y and $\frac{y}{2}$ and $(\frac{y}{2} + 1)$, with $y = 7$ or answer 7		M1M1M1M1A0
	In alt 1 and alt 2, condone missing brackets in equations if not recovered for up to M1M1M1 eg $x + x - 1 + 2x \div 3 = 5$ not recovered		M1M1M1M0A0

Q	Answer	Mark	Comments
11(a)	$\frac{13}{100}$ or 0.13 or 13%	B1	oe fraction, decimal or percentage

Q	Answer	Mark	Comments
11(b)	$\frac{59}{100}$ or 0.59 or 59%	B1	oe fraction, decimal or percentage SC1 answers 13 in (a) and 59 in (b) or $\frac{13}{x}$ in (a) and $\frac{59}{x}$ in (b) where x is an integer ≥ 59

Q	Answer	Mark	Comments
11(c)	$\frac{89}{100}$ or 0.89 or 89%	B1	oe fraction, decimal or percentage SC1 answers 13 in (a) and 89 in (c) or $\frac{13}{x}$ in (a) and $\frac{89}{x}$ in (c), where x is an integer ≥ 89 or answers 59 in (b) and 89 in (c) or $\frac{59}{x}$ in (b) and $\frac{89}{x}$ in (c), where x is an integer ≥ 89
Additional Guidance			
13 in (a) and 59 in (b) and 89 in (c) scores 0, SC1, SC1			

Q	Answer	Mark	Comments
12(a)	$1 \leq a < 10$	B1	allow 1.0 etc
Additional Guidance			
Accept 9.9 for 10			

Q	Answer	Mark	Comments
12(b)	0.0072	B2	B1 7.2×10^3 or 7.2×10^{-3} ignore extra 0s which don't affect the value
Additional Guidance			
0.0072 in working with 7.2×10^{-3} on the answer line			B1

Q	Answer	Mark	Comments
13(a)	$(y =) ax + b$ and $(y =) ax + 2a + b$	B2	any letter for x other than a or b or y B1 $(y =) ax + b$ or $(y =) a(x + 2) + b$ or $(y =) ax + 2a + b$ or substitution of two values for x with a difference of 2 and correct working to show that the output increases by $2a$ eg substituting $x = 3$ and $x = 5$ to get $3a + b$ and $5a + b$
Additional Guidance			
Allow xa for ax throughout			
Do not allow $a \times x + b$ for $ax + b$ unless recovered			
Allow, eg $(x + 2) \times a + b$ for $a(x + 2) + b$			
Do not allow missing brackets unless recovered eg do not allow $x + 2 \times a$ for $a(x + 2)$			
Do not accept written answers without the necessary algebra eg The input has increased by 2 and will then be multiplied by a , so the output will increase by $2a$			B0
Ignore further non-contradictory work if B2 awarded			

Q	Answer	Mark	Comments
13(b)	Alternative method 1 – using k		
	$\frac{f(6)}{f(2)} \left(= \frac{36k}{4k} \right) = 9$ <p>or</p> $f(3) = 9k$	M1	condone eg $k36$
	$\frac{f(6)}{f(2)} = 9 \text{ and } f(3) = 9k$ <p>and No</p>	A1	condone $k9$
	Alternative method 2 – substituting a value for k		
	Identifies a value of k other than 1 and correctly evaluates $\frac{f(6)}{f(2)}$ or $f(3)$	M1	eg $k = 2$ and $\frac{f(6)}{f(2)} = 9$ or $f(3) = 18$
	Identifies a value of k other than 1 and correctly evaluates $\frac{f(6)}{f(2)}$ and $f(3)$ and No	A1	eg $k = 2$ and $\frac{f(6)}{f(2)} = 9$ and $f(3) = 18$ and No
	Additional Guidance		
	$9k$ from $\frac{f(6)}{f(2)}$ is M0, but M1 can be awarded if accompanied by $f(3) = 9k$		
	Do not allow 9 from $\frac{36}{4}$ (unless $\frac{36}{4}$ is from $\frac{36k}{4k}$)		
	Do not allow 9 from $\frac{36k^2}{4k^2}$		
	Students may correctly state that $\frac{f(6)}{f(2)}$ and $f(3)$ are (only) equal when $k = 1$ This may replace 'No' in their answer, but does not score without $9k$ and 9		
	Do not allow unprocessed values, eg 6^2 , 2^2 or 3^2		

Q	Answer	Mark	Comments
14	12 24 30 41	B2	<p>B1</p> <p>their median = $2 \times$ their LQ with the first eight values in order and their UQ and their last number \geq their median</p> <p>or</p> <p>their UQ = $2.5 \times$ their LQ with the first ten numbers in order and their last number \geq their UQ</p> <p>or</p> <p>their range = $2 \times$ their interquartile range with all values in order</p>
Additional Guidance			
Take the boxes to be the LQ, median, UQ and highest value in that order			
Decimal values can score up to B1 eg 11.5 23 29 40 has median = $2 \times$ LQ			B1
Ignore blank boxes for B1			
If all boxes are blank, mark the working lines			

Q	Answer	Mark	Comments
15	True Not true Not true True	B4	B1 each correct answer
Additional Guidance			
Allow a cross if it's the only answer in that row			
If one tick and one or two crosses are given in a row, mark the tick			

Q	Answer	Mark	Comments
Alternative method 1 – equates coefficients and eliminates an unknown			
	$8x - 20y = 52$ and $15x + 20y = 40$ or $6x - 15y = 39$ and $6x + 8y = 16$	M1	oe equates coefficients of one unknown allow one term error
	$8x + 15x = 52 + 40$ or $23x = 92$ or $-15y - 8y = 39 - 16$ or $-23y = 23$	M1dep	oe eliminates an unknown must be correct for their equations
	$x = 4$ and $y = -1$	A2	A1 $x = 4$ from correct method or $y = -1$ from correct method
Alternative method 2 – substitutes for x			
16	$x = 6.5 + 2.5y$ or $x = \frac{8}{3} - \frac{4}{3}y$	M1	oe makes x the subject of one equation allow one term error
	$3(6.5 + 2.5y) + 4y = 8$ or $11.5y = -11.5$ or $2\left(\frac{8}{3} - \frac{4}{3}y\right) - 5y = 13$ or $-\frac{23}{3}y = \frac{23}{3}$	M1dep	oe eliminates x must be correct for their rearrangement
	$x = 4$ and $y = -1$	A2	A1 $y = -1$ from this method

Question 16 continues on the next page

16 cont	Alternative method 3 – substitutes for y		
	$y = 0.4x - 2.6$ or $y = 2 - 0.75x$	M1	oe makes y the subject of one equation allow one term error
	$3x + 4(0.4x - 2.6) = 8$ or $4.6x = 18.4$ or $2x - 5(2 - 0.75x) = 13$ or $5.75x = 23$	M1dep	oe eliminates y must be correct for their rearrangement
	$x = 4$ and $y = -1$	A2	A1 $x = 4$ from this method
	Alternative method 4 – makes the same unknown the subject in both equations		
	$x = 6.5 + 2.5y$ or $x = \frac{8}{3} - \frac{4}{3}y$ or $y = 0.4x - 2.6$ or $y = 2 - 0.75x$	M1	oe makes y or x the subject of one equation allow one term error
	$6.5 + 2.5y = \frac{8}{3} - \frac{4}{3}y$ or $\frac{23}{6}y = -\frac{23}{6}$ or $0.4x - 2.6 = 2 - 0.75x$ or $1.15x = 4.6$	M1dep	oe makes y or x the subject of both equations (maximum one term error) and eliminates y or x must be correct for their rearrangements
	$x = 4$ and $y = -1$	A2	A1 $x = 4$ from correct method or $y = -1$ from correct method
	Additional Guidance		
	Up to M2 may be awarded for correct work seen in multiple attempts, even if not subsequently used		
	In alts 2, 3 and 4 allow rounding or truncating to 1dp or better for up to M1M1 eg (Alt 4) $6.5 + 2.5y = 2.7 - 1.3y$		M1M1
	Answers from trial and improvement or with no working score 0 or 4		

Q	Answer	Mark	Comments
17	Alternative method 1 – expressions in x		
	4 $\pi x^2 \div 2$ or 2 πx^2 or πx^2 or $\pi(3x)^2$ or 9 πx^2 or $2 \times \pi(3x)^2$ or 18 πx^2 or 2 $\pi x(3x)$ or 6 πx^2	M1	oe area of curved face of hemisphere oe area of flat face of hemisphere oe area of one flat face of cylinder oe area of both flat faces of cylinder oe area of curved face of cylinder
	4 $\pi x^2 \div 2 + \pi x^2$ or 3 πx^2 or $\pi(3x)^2 + \pi(3x)^2 + 2\pi x(3x)$ or 9 $\pi x^2 + 9\pi x^2 + 6\pi x^2$ or 24 πx^2	M1dep	oe total surface area of the hemisphere oe total surface area of the cylinder
	3 πx^2 and 24 πx^2 and 1 : 8		either order
	Alternative method 2 – substituting a value for x		
	Substitutes a value for x and works out the area of at least one of area of curved face of hemisphere area of flat face of hemisphere area of one flat face of cylinder area of both flat faces of cylinder area of curved face of cylinder	M1	eg using $x = 5$, at least one of 50 π 25 π 225 π 450 π 150 π
	Substitutes a value for x and works out an expression for the total surface area of the hemisphere or the cylinder		eg using $x = 5$ total surface area of hemisphere = 25 $\pi + 50\pi$ or 75 π or total surface area of cylinder = 225 $\pi + 225\pi + 150\pi$ or 600 π
	Both correct total surface areas for their value of x and 1 : 8	A1	either order

Question 17 continues on the next page

		Additional Guidance	
		1 : 8 or 8 : 1 without correct working or values	
17 cont	Condone π missing consistently for all marks		
	Allow 'correct' and consistent values of π throughout (eg 3, 3.14, $\frac{22}{7}$)		
	Condone use of r for x throughout		
	Do not allow $3\pi x^2$ from $3x \times \pi \times x$ oe		

Q	Answer	Mark	Comments
18	290	B1	

Q	Answer	Mark	Comments			
	$4 \times 3 \times 2 (\times 1) \times 2$ or $5 \times 4 \times 3 \times 2 (\times 1) \times \frac{2}{5}$ or $120 \times \frac{2}{5}$	M1	oe			
19	48					
	SC1 12 or 24 or 72 or 120					
	Additional Guidance					
	12 is the number of possible 5-digit numbers ending in two odd digits					
	24 is the number of possible 5-digit numbers ending in 7 or the number of possible 5-digit numbers ending in 9					
72 is the number of possible 5-digit even numbers						
120 is the number of possible 5-digit numbers						
Ignore any listing of possible numbers						

Q	Answer	Mark	Comments
20	Alternative method 1 – finds K in terms of L and substitutes		
	3K = 4L or K = L + 2M	M1	oe correct equation eg $K = \frac{4L}{3}$ or $L = \frac{3K}{4}$ may be implied by values on diagram
	$\frac{4L}{3} = L + 2M$	M1dep	oe correct equation in L and M eg $4L = 3L + 6M$
	6	A1	condone 6M (= L)
	Alternative method 2 – finds two variables in terms of the other variable		
	Finds one variable in terms of one other eg L is $\frac{3}{4}$ of K	M1	oe fractions, decimals, percentages or ratio eg $K : L = 1 : \frac{3}{4}$ may be implied by values on diagram
	Finds two variables in terms of the other eg L is $\frac{3}{4}$ of K and M is $\frac{1}{8}$ of K	M1dep	oe fractions, decimals, percentages or ratio eg $K : L : M = 1 : \frac{3}{4} : \frac{1}{8}$ may be implied by values on diagram
	6	A1	condone 6M (= L)
	Alternative method 3 – assumes a mass for one unknown		
	Assumes a mass for one unknown and works out the mass of one other	M1	eg $K = 2\text{kg}$ and $L = 1.5\text{kg}$
	Assumes a mass for one unknown and works out the masses of the other two	M1dep	eg $K = 2\text{kg}$ and $L = 1.5\text{kg}$ and $M = 0.25\text{kg}$
	6	A1	condone 6M (= L)
	Additional Guidance		
	Condone 1.33 or better for $\frac{4}{3}$, but 0.125 for $\frac{1}{8}$ must be correct		
	3K : 4L is not enough for M1		
	Ignore units		

Q	Answer	Mark	Comments
21	$(x - 3)^2 - 24$ or $a = 3$ and $b = 24$	B2	B1 $(x - 3)^2 \dots$ or $(x - 3)(x - 3) \dots$ or $a = 3$ (implied by 3, -24) or $x^2 - 2ax + a^2 - b$ or $-2a = -6$ or $2a = 6$ or $a^2 - b = -15$ or correct b for their a
Additional Guidance			
$(x + 3)^2 - 24$ (24 is correct for $a = -3$)			B1
$(x - 6)^2 - 51$ (51 is correct for $a = 6$)			B1
$(x + 6)^2 - 51$ (51 is correct for $a = -6$)			B1

Q	Answer	Mark	Comments
	$a + b \rightarrow 4\sqrt{2}$ $ab \rightarrow 6$ $\frac{b}{a} \rightarrow 3$	B3	B1 each correct match
22	Additional Guidance		
	Two lines from a left-hand box is choice		
	B3		
	B0		

Q	Answer	Mark	Comments
23	Alternative method 1 – subtracting powers of 10 algebraically		
	Denotes the given recurring decimal by a letter and multiplies by one of 10, 100, etc	M1	eg $10x = 1.33\dots$ or $100x = 13.3\dots$
	Denotes the given recurring decimal by a letter and multiplies by one or two of 10, 100, etc and subtracts accordingly	M1dep	eg $10x - x = 1.333\dots - 0.133\dots$ or $9x = 1.2$ or $\frac{1.2}{9}$ or $100x - x = 13.333\dots - 0.133\dots$ or $99x = 13.2$ or $\frac{13.2}{99}$ or $100x - 10x = 13.333\dots - 1.333\dots$ or $90x = 12$ or $\frac{12}{90}$
	$\frac{2}{15}$	A1	
	Alternative method 2 – subtracting powers of 10 numerically		
	Multiplies the given decimal by one of 10, 100, etc	M1	eg $0.\dot{1}3 \times 10 = 1.\dot{3}$
	Multiplies the given decimal by one or two of 10, 100, etc and subtracts appropriately in fraction form	M1dep	eg $0.\dot{1}3 \times 100 = 13.\dot{3}$ and $0.\dot{1}3 \times 10 = 1.\dot{3}$ and $\frac{13.3 - 1.3}{100 - 10}$ or $\frac{12}{90}$
	$\frac{2}{15}$	A1	

Question 23 continues on the next page

23 cont	Alternative method 3 – splitting into a known fraction and a recurring decimal		
	Splits into 0.1 and $0.\overline{03}$ and uses a correct first step from alt 1 or alt 2 with $0.\overline{03}$	M1	eg $10x = 0.33\dots$ or $0.\overline{03} \times 100 = 3.33\dots$ 0.1 does not need to be seen separately at this stage
	Correct method to evaluate $0.\overline{03}$ as a fraction and addition to $\frac{1}{10}$ or $\frac{1}{30} + \frac{1}{10}$ or $\frac{4}{30}$	M1dep	oe fraction
	$\frac{2}{15}$	A1	
	Additional Guidance		
Condone decimals within fractions up to M2 eg $\frac{1.2}{9}$			M2
Equals signs may be implied throughout			
Subtraction signs must be seen or the results correct			
Recurring decimals should be denoted by correct notation or at least two of the recurring digits followed by at least two dots. However, condone missing dots if the result is, or would be, correct eg condone $13.3 - 1.3 = 100x - 10x$			

Q	Answer	Mark	Comments
24	Alternative method 1 – using the equations of the lines $\frac{22-y}{8-0} = 2$ or $22 = 2 \times 8 + c$ or $(c =) 22 - 2 \times 8$ or $c = 6$ or P is at $(0, 6)$ or $(PR =) y = 2x + 6$ or y -coordinate of P is 6 or y -coordinate of Q is 6	M1	oe equation using any letter y is the y -coordinate of P ignore missing brackets may be seen on diagram may be seen on diagram
	$2m = -1$ or $(m =) -\frac{1}{2}$	M1	oe gradient of RQ
	$22 = \text{their } -\frac{1}{2} \times 8 + c$ or $22 = -4 + c$ or $c = 26$ or $(RQ =) y = -\frac{1}{2}x + 26$	M1dep	oe equation in c dep on previous mark oe equation of RQ
	their $(-\frac{1}{2}x + 26) = \text{their } 6$ or x -coordinate of Q is 40	M1dep	oe equation in x where x is the x -coordinate of Q dep on M3 $-\frac{1}{2} = \frac{22 - \text{their } 6}{8 - x}$ implies M4 if their 6 is correct or from correct working
	(40, 6)	A1	

Question 24 continues on the next page

24 cont	Alternative method 2 – using similar triangles		
	Drops a perpendicular from R to point S on PQ and uses $RS = 2PS = 16$ to work out that P is at $(0, 6)$	M1	any or no letter eg $22 - 2 \times 8$
	$2m = -1$ or $(m =) -\frac{1}{2}$ or $\frac{RS}{SQ} = \frac{1}{2}$	M1	oe gradient of RQ
	16×2 or 32	M1dep	length of SQ may be seen on diagram dep on previous mark
	8 + their 32 or x -coordinate of Q is 40	M1dep	
	(40, 6)	A1	
Additional Guidance			
	Note that 40 (for the x -coordinate of Q) implies M3 (on alt 2) and implies M4 if 6 is also seen (on alt 1)		

Q	Answer	Mark	Comments
25	$\sin 30 = \frac{1}{2}$ or $\tan 45 = 1$ or $\cos 30 = \frac{\sqrt{3}}{2}$	M1	oe eg $\tan 45 = \frac{\sqrt{2}}{\sqrt{2}}$ or $4 \sin 30 = 2$ or $2 \cos 30 = \sqrt{3}$ implied by position in the expression may be seen in a table
	substitution of all three correct values		eg $\frac{4 \times \frac{1}{2} - 1}{2 \times \frac{\sqrt{3}}{2}} \text{ or } \frac{2 - 1}{2 \times \frac{\sqrt{3}}{2}} \text{ or } \frac{2 - 1}{\sqrt{3}}$
	$\frac{1}{\sqrt{3}}$ or $\frac{\sqrt{3}}{3}$ $(\frac{1}{\sqrt{3}} \text{ or } \frac{\sqrt{3}}{3} =) \tan 30$ or $x = 30$ with full working seen for M3	M1dep	
			Additional Guidance
	Allow $\sqrt{1}$ for 1 throughout		
	Reference to 30° being an acute angle is not required		

Q	Answer	Mark	Comments
Alternative method 1			
20π ÷ 2π or 10		M1	oe may be seen on diagram implied by diameter = 20
$x^2 + x^2 = (\text{their } 10)^2$ or $2x^2 = 100$ or $x^2 = 50$ or their $10 \times \sin 45$ or their $10 \times \cos 45$ or their $10 \times \frac{1}{\sqrt{2}}$		M1	oe any letter (condone a) their 10 is their length OQ (the radius of the circle)
$\sqrt{\text{their } 10^2 \div 2}$ or $\sqrt{50}$ or $5\sqrt{2}$ or $4 \times \sqrt{50}$ or $4 \times \text{their } 10 \times \sin 45$ or $4 \times \text{their } 10 \times \cos 45$ or $40 \times \frac{1}{\sqrt{2}}$ or $\frac{40\sqrt{2}}{2}$ or $20\sqrt{2}$		M1dep	oe value for the length of one side of the square or the perimeter of the square eg $\frac{10}{\sqrt{2}}$ dep on previous mark
2 with full working seen for M3		A1	

Question 26 continues on the next page

26 cont	Alternative method 2		
	20π ÷ 2π or 10 or side length of square = $5\sqrt{a}$	M1	oe may be seen on diagram implied by diameter = 20
	(Perimeter of square = $20\sqrt{a}$ and) side length of square = $5\sqrt{a}$ and $(5\sqrt{a})^2 + (5\sqrt{a})^2 = (\text{their } 10)^2$	M1	oe their 10 is their length OQ (the radius of the circle) condone missing brackets if recovered
	25a + 25a = (their 10) ² or 50a = 100	M1dep	dep on M1M1
	2 with full working seen for M3	A1	
	Additional Guidance		
	2 with no working		M0M0M0A0
	$\sqrt{2}$ on answer line (may score method marks)		A0

Q	Answer	Mark	Comments
27	(Total time =) $\frac{30}{a} + \frac{30}{b}$	M1	oe eg $\frac{30b}{ab} + \frac{30a}{ab}$ or $\frac{30(b+a)}{ab}$
	correct expression for total distance \div total time	M1dep	eg $(30 + 30) \div \left(\frac{30}{a} + \frac{30}{b} \right)$ or $60 \div \frac{30(b+a)}{ab}$ or $60 \times \frac{ab}{30(b+a)}$
	$60 \times \frac{ab}{30(a+b)} = \frac{2ab}{a+b}$	A1	condone $b+a$ for $a+b$ condone $30a + 30b$ for $30(a+b)$
Additional Guidance			
Students can gain M1M1 if they incorrectly simplify a correct expression for total time before forming the division eg $\frac{30}{a} + \frac{30}{b} = \frac{60}{a+b}$ followed by $60 \div \frac{60}{a+b}$			M1M1A0
Allow correct cancellation of 60 and 30 at any stage of the working			