

International GCSE Maths

Apart from questions 2, 6, 10, 13a, 18ai (where the mark scheme states otherwise) the correct answer, unless clearly obtained by an incorrect method, should be taken to imply a correct method

Q	Working	Answer	Mark	Notes
1	e.g. $36 \div (2 + 6) (= 4.5)$ or $36 \div \frac{2+6}{3+2+6} (= 49.5)$ oe or Asha = £9 OR Julie = £27			M1
	e.g. $3 \times "4.5"$ or " $49.5 \times \frac{3}{3+2+6}$ " or " $9 \times \frac{3}{2}$ " or " $27 \times \frac{3}{6}$ "			M1 or an answer of $\frac{27}{2}$
		13.5(0)	3	A1 SCB1 for $36/5 \times 6 (=43.2)$ or $36/9 \times 2 (=8)$
				Total 3 marks

2	e.g. $\frac{16}{5}$ and $\frac{21}{8}$ oe			M1 both fractions expressed as improper fractions
	e.g. $\frac{16^2}{5} \times \frac{21}{8^1}$ OR $\frac{336}{40}$ oe			M1 correct cancelling OR multiplication of numerators and denominators without cancelling
	e.g. $\frac{16}{5} \times \frac{21}{8} = \frac{336}{40} = \frac{42}{5} = 8\frac{2}{5}$ or $\frac{16}{5} \times \frac{21}{8} = \frac{336}{40} = 8\frac{16}{40} = 8\frac{2}{5}$ or $\frac{16^2}{5} \times \frac{21}{8^1} = \frac{42}{5} = 8\frac{2}{5}$ or candidate clearly shows that in the question, the result of $8\frac{2}{5} = \frac{42}{5}$ and that their answer becomes $\frac{42}{5}$	shown	3	A1 Dep on M2 for conclusion to $8\frac{2}{5}$ from correct working – either sight of the result of the multiplication e.g. $\frac{336}{40}$ must be seen or correct cancelling prior to the multiplication to $\frac{42}{5}$ NB: use of decimals scores no marks
				Total 3 marks

3 a	e.g. $d - g = 2ac$ $\frac{d}{2c} = \frac{g}{2c} + a$			M1 for a correct first step e.g. subtract g from both sides OR divide all terms by 2 OR divide all terms by c OR divide all terms by $2c$
		$a = \frac{d - g}{2c}$	2	A1 oe
b		$3f(3e - 4)$	2	B2 (B1 for $3(3ef - 4f)$ or $f(9e - 12)$ or $3f(ke - 4)$ or $3f(3e - m)$ where $k \neq 0$ and $m \neq 0$)
c	$x^2 - 5x + 2x - 10$			M1 for any 3 correct terms or for 4 out of 4 correct terms ignoring signs or $x^2 - 3x - 10$ or for $\dots - 3x - 10$
		$x^2 - 3x - 10$	2	A1
d	$\frac{n^{11}}{n^5}$ OR $n^{-1} \times n^7$ OR $n^4 \times n^2$ OR $n^4 \times n^7 \times n^{-5}$ OR $n^{(11)} \div n^5 = n^{(11) - 5}$			M1 for simplifying two terms
		n^6	2	A1
				Total 8 marks

4 ai		b, l, u, e, g, r, y	1	B1 No incorrect or repeats
a ii		w, h, i, t	1	B1 No incorrect or repeats
b		No with reason	1	B1 eg 'e is in all three sets' OR 'all three sets share a member' OR $B \cap G \cap W = (\{e\})$
				Total 3 marks

5	$\pi \times 7.2^2 \div 2$ (= 81.4....)			M1 allow 81.3 – 81.5 for area of semi circle
	“81.4” \div 6 (= 13.5...) or 12×6 (= 72) or “81.4” \div 12 (= 6.7...)			M1 (dep) allow 13.5 – 13.6 for the number of boxes needed (NB: $12 \times 6 = 72$ alone is 0 marks)
		No with correct figures	3	A1
				Total 3 marks

6	$(x \pm 9)(x \pm 4)$	$\frac{-(-5) \pm \sqrt{(-5)^2 - 4 \times 1 \times (-36)}}{2 \times 1}$ or $\frac{5 \pm \sqrt{25+144}}{2}$		M1 or $(x + a)(x + b)$ where $ab = -36$ or $a + b = -5$ OR correct substitution into quadratic formula (condone one sign error in a , b or c) (if + rather than \pm shown then award M1 only unless recovered with answers)
	$(x - 9)(x + 4)$	$\frac{5 \pm \sqrt{169}}{2}$ or $\frac{5 \pm 13}{2}$		M1 or $\frac{5 \pm \sqrt{169}}{2}$ or $\frac{5 \pm 13}{2}$
		9, -4	3	A1 dep on at least M1
				Total 3 marks

7	$20.40 \div (1 - 0.15)$			M2 for a complete method eg $20.40 \div (1 - 0.15)$ (M1) for $20.40 \div (100 - 15)$ (= 0.24) or e.g. $0.85x = 20.40$
		24	3	A1
				Total 3 marks

8	$28 \times 5 (= 140)$ OR $26.5 \times 2 (= 53)$			M1 or 87
	$(28 \times 5 - 26.5 \times 2) \div (5 - 2)$			M1 for a complete method
		29	3	A1
				Total 3 marks

9	$1.5 \times 2 \times 8 (= 24 \text{ (cm}^3\text{)})$			M1 for finding the volume of the cuboid
	e.g. $(V =) \frac{5.73 \times 1000}{19.32} (= 296.58\ldots)$ or $(M =) 19.32 \times "24" (= 463.68)$			M2 complete method to find the volume of statue or the mass of one block, could work in g or kg (if not M2 then award M1 for correct use of density formula e.g. $19.32 = \frac{5.73 \times 1000}{V}$ or $19.32 = \frac{M}{"24"}$)
	e.g. "296.58" \div "24" (= 12.3576...) or "5730" \div "463.68" (= 12.3576...)			M1 could work in g or kg
		13	5	A1 cao
				Total 5 marks

10	e.g. $6(x - 1) (= 6x - 6)$			M1 method to find expression for perimeter of hexagon
	e.g. $2(x + 5) + 2x - 3 (= 4x + 7)$			M1 method to find expression for perimeter of triangle
	" $6x - 6$ " = " $4x + 7$ "			M1 (dep on at least M1) for equating both expressions
	e.g. $6x - 4x = 7 + 6$			M1 (dep on previous M1 and equation of the form $ax + b = cx + d$) for rearranging the x terms on one side and the numerical terms on the other and all expansions correct.
		5.5	5	A1 oe (dep on M2)
				Total 5 marks

11	a		4.35	1	B1 accept 4.34 ⁸
	b		4.25	1	B1 cao
c	17.5 or 9.35 or $e - f$ where $17 < e \leq 17.5$ and $9.35 \leq f < 9.4$				M1
			8.15	2	A1
					Total 4 marks

12	a		23	1	B1 accept 22 – 24
	b	e.g. 29 – 17			M1 For subtracting readings from 15 and 45
			12	2	A1 accept 10 – 14
c					B1 ft comparison of the medians
		Two comparisons (at least one of which must be in context)	2		B1 ft comparison of the IQR Note: to award 2 marks at least one comparison must be in context
					Total 5 marks

13	a	<p>e.g. $x = 0.57272\dots$ and $100x = 57.272\dots$ OR e.g. $10x = 5.7272\dots$ and $1000x = 572.72\dots$</p>			M1 For 2 recurring decimals with correct algebraic labels that when subtracted give a whole number or terminating decimal eg 56.7 or 567 etc e.g. $100x = 57.272\dots$ and $x = 0.57272\dots$ OR $1000x = 572.72\dots$ and $10x = 5.7272\dots$ with intention to subtract. (If recurring dots not shown then showing at least the digits 57272, ie 5sf)
		<p>e.g. $100x - x = 57.272\dots - 0.57272\dots = 56.7$ and $\frac{56.7}{99} = \frac{63}{110}$ or $1000x - 10x = 572.72\dots - 5.7272\dots = 567$ and $\frac{567}{990} = \frac{63}{110}$</p>	Shown	2	A1 for completion to $\frac{63}{110}$
	b	$\frac{3}{2-\sqrt{y}} \times \frac{2+\sqrt{y}}{2+\sqrt{y}}$ or $6+3\sqrt{y}$ or $4-y$			M1 for multiplying numerator and denominator by $(2+\sqrt{y})$ or a correct expression for the numerator or denominator
			$\frac{6+3\sqrt{y}}{4-y}$	2	A1
					Total 4 marks

14	$(AOC =) 38 \times 2 (= 76)$		4	M1
		52		A1
				B2 (dep on M1) for all reasons relevant to their method – underlined words must be seen. <u>angle</u> at the <u>centre</u> is $2 \times$ (double) angle at <u>circumference</u> / <u>angle</u> at <u>circumference</u> is $\frac{1}{2}$ <u>angle</u> at <u>centre</u> <u>angles</u> in a <u>triangle</u> add to 180° or angles in a <u>triangle</u> add to <u>180°</u> base angles in an <u>isosceles</u> triangle (are equal)
				If not B2 then award B1 (dep on M1) for a correct circle theorem
				Total 4 marks

15	e.g. $(EF =) 12\cos 40 (= 9.19\dots)$ or $(FD =) 12\sin 40 (= 7.71\dots)$ and $(EF =) \sqrt{12^2 - "7.71"^2} (= 9.19\dots)$			M2 complete method to find EF (if not M2 then M1 for a correct statement involving EF e.g. $\frac{EF}{12} = \cos 40$)
	e.g. $\frac{"9.19"}{EG} = \tan 28$ or $\tan 62 = \frac{EG}{"9.19"}$ or $\frac{"9.19"}{FG} = \sin 28 (= 19.5\dots)$ and $"19.5"^2 - "9.19"^2 (= 298.9\dots)$			M1 (dep on M2) for a correct trig statement involving EG or complete method to find FG and a correct start to Pythagoras process
		17.3	4	A1 accept 17.2 – 17.3
				Total 4 marks

16	$0.42 \div 0.6 (= 0.7)$ oe			M1 (indep)
	$1 - "0.7" (= 0.3)$ oe OR $1 - 0.6 (= 0.4)$ oe			M1 (indep)
	$"0.3" \times "0.4"$ oe OR $1 - (0.42 + 0.6 \times "0.3" + "0.4" \times "0.7")$ oe			M1 for a complete method
		0.12	4	A1 oe
				Total 4 marks

17 a		9	1	B1
b		$f(x) \geq 0$	1	B1 accept $y \geq 0$ or $f \geq 0$
c	$(g(2) =) \frac{4}{2+3} \left(= \frac{4}{5} \right)$ oe			M1 or for sight of $fg(x)$ e.g. $\left(\frac{4}{x+3} - 4 \right)^2$
		10.24	2	A1 oe e.g. $\frac{256}{25}$
				Total 4 marks

18	ai			B1	tangent drawn at $P (x = 2)$
				M1	(dep on B1) for a method to find gradient e.g. <u>difference in y-values</u> <u>difference in x-values</u>
		-0.6	3	A1	(dep on B1) accept answers in range -0.4 to -0.7 and from correct figures for their line
	aii	e.g. $y = -0.6x + c$ or $y = mx + 3.6$ or $2.4 = -0.6 \times 2 + c$		M1	for start of method to find the tangent equation e.g. $y = mx + c$ where m is their gradient or $y = mx + c$ where c is the y -intercept for their tangent or for substituting a point from their tangent e.g. (2, 2.4) into $y = mx + c$ where m is their gradient
		$y = -0.6x + 3.6$	2	A1	ft their gradient from (i) and intercept of their tangent, so long as intercept / value of c is > 3
	b	3		B1	
		-1	2	B1	
					Total 7 marks

19	<p>eg $5 \times 2x + 10 \times x = 160$</p> <p>OR</p> <p>$160 \div 2 (= 80)$ [freq of one bar]</p> <p>OR</p> <p>$40 \times 5 + 20 \times 10 (= 400)$ [total no. of sml squares]</p> <p>OR</p> <p>$160 \div 16 (= 10)$ [students per 1cm^2]</p> <p>OR</p> <p>$1\text{cm}^2 = 10$ students</p> <p>OR</p> <p>e.g. 5 small squares = 2 students oe</p>			M1 for setting up an appropriate equation OR finding the area of the 2 nd or 3 rd bar OR finding the total number of small squares OR for finding the number of students per 1cm^2 or $1\text{cm}^2 = 10$ students OR other appropriate scale e.g. 5 small squares = 2 students
	<p>$'x' = 8$</p> <p>OR</p> <p>8 or 16 seen in the correct position on the vertical scale</p> <p>OR</p> <p>$160 \div "400" (= 0.4 \text{ oe})$</p>			M1 for finding frequency density OR method to find the frequency of the 1 st , 4 th or 5 th bar (1 st is 108, 4 th is 90, 5 th is 12)
	<p>$"7.2" \times 15 + 160 + "6" \times 15 + "2.4" \times 5$</p> <p>OR $160 + "0.4" \times (18 \times 15 + 15 \times 15 + 5 \times 6)$</p>			M1 (dep on at least M1) for a complete method to find the total frequency (allow one error or one repeat but no omission)
		370	4	A1
				Total 4 marks

20	$\frac{1}{3} \times \pi \times r^2 \times 2h \left(= \frac{2}{3} \pi r^2 h \right)$ OR $\frac{1}{3} \times \pi \times (0.5r)^2 \times h \left(= \frac{1}{12} \pi r^2 h \right)$			M1 for finding the volume of the small or large cone
	$"\frac{2}{3} \pi r^2 h" - "\frac{1}{12} \pi r^2 h" \left(= \frac{7}{12} \pi r^2 h \right)$			M1 (dep) method to find the volume of the frustum (condone missing brackets)
	$"\frac{2}{3} \pi r^2 h" - "\frac{1}{12} \pi r^2 h" = \frac{4\pi r^3}{3}$			M1 equating volume of frustum and sphere (must be correct including brackets)
	e.g. $\frac{7}{12} \pi r^2 h = \frac{4\pi r^3}{3}$			M1 for a correct simplified formula (1 term on each side)
		$\frac{7}{16} h$	5	A1 accept $0.4375h$
				Total 5 marks

21	$CB = 13 \sin 40$ ($= 8.3562\dots$)			M1
	$\frac{1}{2} \times 6 \times "8.35\dots" \times \sin ACB = 22$			M1
	Acute version of $ACB = \sin^{-1} \left(\frac{22}{\frac{1}{2} \times 6 \times "8.35\dots"} \right)$ ($= 61.35\dots$)			M1
	$ACB = 180 - "61.353\dots"$ ($= 118.647\dots$)			M1
	$AB^2 = 6^2 + "8.35\dots"^2 - 2 \times 6 \times "8.35\dots" \times \cos "118.64"$ ($= 153.98\dots$)			M1
		12.4	6	A1 accept $12.3 - 12.5$
				Total 6 marks

22 a			2	M1 for one correct value
		2.5, -60	A1 oe e.g. -2.5 & 120	
			SC M1 for drawing cos curve	
bi		(2, 5)	1	B1
bii		(4, -2)	1	B1
				Total 4 marks

23	$(v =) 3t^2 + 2 \times 4t - 5$			M1 2 out of 3 terms differentiated correctly
	$3T^2 + 8T - 5 = V \text{ OR } 3T^2 + 8T - 5 - V = 0$			A1 correct equation
	$3(T^2 + \frac{8}{3}T) - 5$ OR $3(T^2 + \frac{8}{3}T - \frac{5}{3})$	$(T =) \frac{-8 \pm \sqrt{8^2 - 4 \times 3 \times (-5 - V)}}{2 \times 3}$		M1 attempt to complete the square OR use quadratic formula (condone one sign error in a, b or c and ft their quadratic with mistake in a or b) (condone + instead of \pm)
	$\left(T + \frac{4}{3}\right)^2 = \left(\frac{4}{3}\right)^2 + \frac{V+5}{3}$	$(T =) \frac{-8 \pm \sqrt{124+12V}}{6}$		M1 sight of this method mark implies the previous M1 (condone + instead of \pm) (ft their quadratic with mistake in a or b)
	$T = \frac{-4}{3} \pm \frac{1}{3}\sqrt{16+3V+15}$	$(T =) \frac{-8 \pm 2\sqrt{31+3V}}{6}$		M1 (condone + instead of \pm) (ft their quadratic with mistake in a or b)
		$\frac{-4 + \sqrt{31+3V}}{3}$	6	A1 accept $k = 31$ and $m = 3$
				Total 6 marks